



Fund “Nauka” Project № 22008 Resume – Competition-Based Session 2022:

“Physical activity in patients on hemodialysis and opportunities for its optimization”

Project leader: Prof. Valentina Hristova Madjova, MD, PhD

Chronic kidney disease (CKD) is a condition causing irreversible loss of the major renal functions: excretory, regulatory and incretory, leading to a significant disruption of the function of all organs and systems in the body. It is a global health problem associated with reduced life expectancy. In patients with end-stage renal disease (ESRD) on haemodialysis (HD), the life expectancy is reduced by nearly 75% compared to their healthy peers. There is also a limited functional physical capacity, (about 70% of clinically healthy individuals, with values decreasing to 50% with the onset of HD). HD patients also have a high level of disability due to the major complications of CKD and a sedentary lifestyle which negatively affects their ability to perform activities of daily living.

The research objective of this project is to perform a comprehensive study of the levels of physical activity in hemodialysis patients with stratification of risk factors for disability and to evaluate the impact of aerobic intradialytic exercise on functional status and clinical outcome with a view to the prognosis, rehabilitation and quality of life of the ESRD patients.

In order to accomplish the set goal, the following tasks are differentiated:

1. To study the impact of chronic dialysis treatment on physical capacity and individual quality of life of patients;
2. To identify the main causes of deterioration of physical capacity and risk stratification;
3. To evaluate the impact of specialized intradialytic exercises on clinical, functional status and individual quality of life of chronic dialysis patients.

The subjects of the study are 80 HD patients randomized into work and control groups.

The research methods to be used are: documentary, clinical, sociological, statistical and laboratory.

As a result of this study, it is expected to create and implement into routine clinical practice an original aerobic intradialytic exercise algorithm for improvement of prognosis, better clinical outcome and quality of life of HD patients.

Expected results:

1. ESRD patients have low levels of physical activity. To identify the causes of low physical activity in these patients and to develop strategies to address them.
2. To validate the hypothesis that intradialytic cardio-velo exercise training is appropriate for positively influencing functional status, physical and emotional health, and individual quality of life of patients on dialysis treatment.

3. Intradialytic exercise positively influences patients' attitudes towards exercise.
4. To establish a clinically applicable methodology for intradialytic exercise at reduced financial cost and improved patient outcome and prognosis.

The main activity of the project was the first study in Bulgaria to apply aerobic intradialysis exercises with a specialized wheel in patients on chronic dialysis. CKD is a serious global health problem associated with reduced life expectancy. In patients with ESKD undergoing HD survival is about 75% lower than that of their peers. There is also a significantly reduced physical functional capacity and disability. The aim of this dissertation is to prove that kinesitherapy improves motor activity, social rehabilitation, and QoL in patients with ESKD. The selection of the studied individuals was made among the patients with CKD who underwent treatment in 2023-2024 at the dialysis center at the UMHAT "St. Marina" Varna. 81 individuals were included in the sample, representing 45% of those who underwent hemodialysis treatment for the studied period. Detailed history and physical status were taken for all participants. The following examinations were performed and completed: Laboratory tests, research for functional status, quality of life assessment. The experimental group's protocol included an aerobic exercise program using a bicycle ergometer – cycling at the beginning of the dialysis procedure with low intensity of load. The control group did not perform intradialytic exercises. As a result of the project activity, an algorithm was developed for assessing and increasing the physical activity of patients on dialysis treatment with application in controlled hospital/outpatient conditions. The project activity is the implementation of tertiary prophylaxis aimed at preventing the development of complications associated with the end-stage of CKD, maintaining maximum functional capacity and improving the quality of life of these patients. As a result of the project activity, it was found that in patients performing aerobic intradialysis exercises with a specialized wheel, their functional status is affected. This gives us reason to recommend that these exercises be used for rehabilitation. The application of renal rehabilitation in the future is one of the factors that will improve the quality of life in patients with CKD on dialysis treatment.